

SI:

Math 122 Test 1

February 3, 2026

EF:

Name_____

1	
2	
3	
4	
5	
Total	

Directions:

1. No books, notes or temperatures above freezing. You may use a calculator to do routine arithmetic computations. You may *not* use your calculator to store notes or formulas. You may not share a calculator with anyone.
2. You should show your work, and explain how you arrived at your answers. A correct answer with no work shown (except on problems which are completely trivial) will receive no credit. If you are not sure whether you have written enough, please ask.
3. You may not make more than one attempt at a problem. If you make several attempts, you must indicate which one you want counted, or you will be penalized.
4. You may leave as soon as you are finished, but once you leave the exam, you may not make any changes to your exam.

1. (20 points)

(a) Compute $\int \frac{2x}{x^4 + 2x^2 + 1} dx$

(b) Compute $\int \frac{1 - \ln x}{x^2} dx$

2. (20 points)

(a) Compute $\int \cos^5 x \, dx$

(b) Compute $\int \frac{1}{\cos^4 x} \, dx$

3. (20 points)

(a) Compute $\int \frac{1}{\cosh^4 x} dx$

(b) (10 points) $\int \frac{6x^2 + 2x - 2}{x^3 - x} dx$

4. (20 points)

(a) Compute $\int \frac{1}{\sqrt{x^2 + 4}} dx$

(b) Compute $\int \frac{1}{x\sqrt{4 - (\ln x)^2}} dx$

5. (20 points)

(a) Compute $\int_0^4 \frac{2x}{x^2 - 9} dx$

(b) Use the chart below and the Trapezoidal method with $n = 4$ sub-intervals to approximate $\ln 2$, where

$$\ln 2 = \int_1^2 \frac{1}{x} dx$$

x	$f(x)$	M.F.	P

FORMULA PAGE

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\sin 2x = 2 \sin x \cos x$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

$$(\tan x)' = \sec^2 x$$

$$(\sec x)' = \sec x \tan x$$

$$(\csc x)' = -\csc x \cot x$$

$$(\cot x)' = -\csc^2 x$$

$$(e^x)' = e^x$$

$$(\ln x)' = \frac{1}{x}$$

$$(\arcsin x)' = \frac{1}{\sqrt{1 - x^2}}$$

$$(\arctan x)' = \frac{1}{1 + x^2}$$

$$(\operatorname{arcsec} x)' = \frac{1}{|x|\sqrt{x^2 - 1}}$$

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

$$\cosh^2 x - \sinh^2 x = 1$$

$$(\sinh x)' = \cosh x$$

$$(\cosh x)' = \sinh x$$

$$(\operatorname{arcsinh} x)' = \frac{1}{\sqrt{x^2 + 1}}$$

$$(\operatorname{arccosh} x)' = \frac{1}{\sqrt{x^2 - 1}}$$

$$(\operatorname{arctanh} x)' = \frac{1}{1 - x^2}$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(c) = \frac{1}{b - a} \int_a^b f(x) dx$$

$$\int \sec x \, dx = \ln |\sec x + \tan x| + C$$

$$\int \sec^3 x \, dx = \frac{1}{2} [\sec x \tan x + \ln |\sec x + \tan x|] + C$$

$$\int \csc x \, dx = \ln |\csc x - \cot x| + C$$

$$1 + 1 = 2$$